

Durham Research Online

Deposited in DRO:

17 July 2014

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Innes, J.B. and Zong, Y. and Wang, Z. and Chen, Z. (2014) 'Climatic and palaeoecological changes during the mid- to Late Holocene transition in eastern China : high-resolution pollen and non-pollen palynomorph analysis at Pingwang, Yangtze coastal lowlands.', *Quaternary science reviews.*, 99 . pp. 164-175.

Further information on publisher's website:

<http://dx.doi.org/10.1016/j.quascirev.2014.06.013>

Publisher's copyright statement:

NOTICE: this is the author's version of a work that was accepted for publication in *Quaternary Science Reviews*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Quaternary Science Reviews*, 99, 2014, 10.1016/j.quascirev.2014.06.013.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Manuscript Number: JQSR-D-13-00387R2

Title: Climatic and palaeoecological changes during the mid- to Late Holocene transition in eastern China: high-resolution pollen and non-pollen palynomorph analysis at Pingwang, Yangtze coastal lowlands.

Article Type: Research and Review Paper

Keywords: Neolithic; coastal east China; palynology; climate change; Neoglacial; vegetation history

Corresponding Author: Dr. James Innes,

Corresponding Author's Institution: Durham University

First Author: James Innes

Order of Authors: James Innes; Yongqiang Zong; Zhanghua Wang; Zhongyuan Chen

Abstract: The transition to the Late Holocene/Neoglacial occurred as a worldwide process of climatic deterioration from the optimum thermal conditions of the mid-Holocene, culminating in an abrupt decline around 4200 cal yr ago, in a period of severe climatic deterioration that lasted for two or three centuries. This sudden climatic event has been recorded in many proxy data archives from around the world, and its effects were manifest in different ways depending on the reaction of regional weather systems and conditions, but often as greatly increased aridity and/or cold temperatures. It has been regarded as causing or contributing to the sudden collapse of several well-established human societies at that time, including advanced agricultural Late Neolithic cultures in eastern China. We have used high-resolution pollen and non-pollen palynomorph analysis to examine the nature of this climatic transition through its impacts on the vegetation and hydrology at Pingwang, a site in the Yangtze coastal lowlands which has no evidence of complicating environmental influences such as sea-level rise or significant human land-use activity, factors previously suggested as alternative reasons for changes in forest composition. Our results show two phases of forest alteration, one gradual from about 5500 cal BP and one sudden at about 4200 cal BP, in which the frequencies of subtropical forest elements fall and are replaced by those of conifers and cold-tolerant trees. Total arboreal pollen frequencies do not decline and the proportion of temperate forest trees, tolerant of a wide range of temperatures, remains unchanged throughout, both ruling out human land clearance as a cause of the change in forest composition. As these dates accord very well with the known timings of climate deterioration established from other proxy archives in the region, we conclude that climate was the main driver of vegetation change in eastern China at the mid- to Late Holocene transition. Our hydrological results support the view that a combination of rising local water level and climatic cooling during the 4200 cal BP event was the probable cause of societal collapse in the lower Yangtze valley.

Climatic and palaeoecological changes during the mid- to Late Holocene transition in eastern China: high-resolution pollen and non-pollen palynomorph analysis at Pingwang, Yangtze coastal lowlands.

James B. Innes ^{a,*}, Yongqiang Zong ^b, Zhanghua Wang ^c, Zhongyuan Chen ^c

^aGeography Department, Durham University, South Road, Durham, DH1 3LE, UK

^bDepartment of Earth Sciences, The University of Hong Kong, Hong Kong SAR, China

^cDepartment of Geography, East China Normal University, Shanghai 200062, China

Abstract:

The transition to the Late Holocene/Neoglacial occurred as a worldwide process of climatic deterioration from the optimum thermal conditions of the mid-Holocene, culminating in an abrupt decline around 4200 cal yr ago, in a period of severe climatic deterioration that lasted for two or three centuries. This sudden climatic event has been recorded in many proxy data archives from around the world, and its effects were manifest in different ways depending on the reaction of regional weather systems and conditions, but often as greatly increased aridity and/or cold temperatures. It has been regarded as causing or contributing to the sudden collapse of several well-established human societies at that time, including advanced agricultural Late Neolithic cultures in eastern China. We have used high-resolution pollen and non-pollen palynomorph analysis to examine the nature of this climatic transition through its impacts on the vegetation and hydrology at Pingwang, a site in the Yangtze coastal lowlands which has no evidence of

complicating environmental influences such as sea-level rise or significant human land-use activity, factors previously suggested as alternative reasons for changes in forest composition. Our results show two phases of forest alteration, one gradual from about 5500 cal BP and one sudden at about 4200 cal BP., in which the frequencies of subtropical forest elements fall and are replaced by those of conifers and cold-tolerant trees. Total arboreal pollen frequencies do not decline and the proportion of temperate forest trees, tolerant of a wide range of temperatures, remains unchanged throughout, both ruling out human land clearance as a cause of the change in forest composition. As these dates accord very well with the known timings of climate deterioration established from other proxy archives in the region, we conclude that climate was the main driver of vegetation change in eastern China at the mid- to Late Holocene transition. Our hydrological results support the view that a combination of rising local water level and climatic cooling during the 4200 cal BP event was the probable cause of societal collapse in the lower Yangtze valley.

KEYWORDS: Neolithic; coastal east China; palynology; climate change; Neoglacial; vegetation history

*Corresponding author (e-mail address: j.b.innes@durham.ac.uk)

1. Introduction

Holocene climate history is now relatively well understood at the global scale, with the recognition of a series of significant temperature events that occurred within the longer-term evolution of the present interglacial's temperate climate, some of which represent tipping points, major shifts that mark phase transitions between longer periods of more stable thermal conditions. One of the most important of these climatic shifts is Holocene event 3 (Bond et al., 2001), occurring relatively abruptly around 4200 years ago (hereafter cal BP) and having major environmental impacts worldwide, recognised (Walker et al., 2012) as the global transition from early and mid-Holocene thermal maxima (Renssen et al., 2012) to the Late Holocene (Neoglacial), with its colder, more extreme and more variable climatic regimes (Jessen et al., 2005; Marchant and Hooghiemstra, 2004; Wanner et al., 2011). This major climatic shift to unstable Neoglacial conditions in the centuries preceding 4000 cal BP has been recorded across the globe in a range of proxy climate data archives (Gear and Huntley, 1991; Jian et al., 1996; Phadtare, 2000; Stott et al., 2004; Booth et al., 2005; Magny et al., 2009; Geirsdóttir et al., 2013), affecting major atmospheric systems such as the North Atlantic Oscillation (Olsen et al. 2012), ENSO (Schulmeister and Lees, 1995), the Indian Monsoon (Overpeck et al., 1996; Gupta et al., 2003; Staubwasser et al. 2003) and the East Asian Monsoon (EAM) through global climatic teleconnections (Wang et al., 2000; Liu et al., 2004; Hong et al., 2005; Tan et al., 2008). A gradual onset of globally cooler conditions can be noted at around 5500 cal BP, with the major Neoglacial intensification at ca 4200 cal BP (Geirsdóttir et al., 2013). The EAM, which forms the focus of this paper, accords very well with the global data, with ample evidence of its progressive weakening after the mid-Holocene at ca 5500 cal BP, culminating in an abrupt decrease in its strength in the centuries before 4000 cal BP (An, 2000; An et al., 2000; Morrill et

al., 2003; He et al., 2004; Yuan et al, 2004; Wang et al., 2005; Selveraj et al., 2007; Cosford et al., 2008; Cai *et al.*, 2010).

1.1. Neoglacial climatic deterioration and human societies

The link between climatic and societal change can be very strong (Perry and Hsu, 2000; Caseldine and Turney, 2010; Zhang et al., 2011). It is not deterministic (Coombes and Barber, 2005) to assume that the change from the mid-Holocene climatic optimum to the much less congenial Neoglacial climate would have had significant effects upon human societies, particularly those that had developed intensive agricultural systems that supported high, sedentary populations, but which had become dependent upon stable, favourable and reliable climatic conditions. The more economically and socially specialized such societies became, the more vulnerable they would have been to any rapid environmental change (O’Sullivan, 2008). Human communities are resilient and adaptable (Anderson et al., 2007; Lu, 2007) and were able to cope and even flourish during the gradual climatic decline from the mid-Holocene thermal maximum around the world, the development of advanced agrarian societies based on the control of water resources in the major river valleys of the Middle East and China being good examples. In China such advanced, highly productive farming systems sustained the dense Neolithic populations of the Longshan, Shijiahe, Qijia and Liangzhu cultures in the Yellow River and Yangtze valleys during the millennium after 5000 cal BP (Liu and Feng, 2012; Zhuang et al., 2014). Abrupt climatic deterioration would have been difficult for such complex agrarian societies to cope with, however (Mo et al., 2011), and there is abundant archaeological and palaeoecological evidence that the rapid climate changes around 4200 cal BP caused severe economic and political dislocation, and even societal collapse, in many regions of the world (Dalfes et al., 1997; Sandweiss et al., 1999; Peiser, 1998; deMenocal, 2001). Major civilizations

disintegrated at this time in India and the Middle East and the coincidence of societal failure and settlement abandonment with the rapid change to Neoglacial cold and arid conditions in these areas implies an environmental cause and a cultural effect, with climate as the driving force (Weiss et al., 1993; Cullen et al., 2000; Staubwasser et al., 2003; Drysdale et al., 2006; Staubwasser and Weiss, 2006; Riehl, 2012), although of course such a direct relationship can never be proven conclusively.

1.2. The East China example

One of the world's regions that has a clear correlation between cultural 'collapse' and the 4200 cal BP climate event is eastern China, and in particular the lower Yangtze valley, one of the historic heartlands of Chinese society and agricultural development. Liu and Feng (2012) and Wu et al. (2012) have evaluated the evidence for the development in the fifth millennium cal BP of the advanced agrarian cultures of central and eastern China mentioned above, and their sudden, almost synchronous demise around 4200 cal BP, when severe climate deterioration occurred. Many major Neolithic archaeological sites have culturally sterile sediments of this age that seal the rich cultural layers and are interpreted as flood deposits. Many authors (e.g. Yu et al., 2000, 2003; Jin and Liu, 2002; Wu and Liu, 2004; Zhang et al., 2005; Gao et al., 2007; Chen et al., 2008) are convinced that the 4200 cal BP event must have been responsible for site abandonment and for the culturally impoverished interlude of a few centuries recorded almost everywhere in north and east China around 4200 cal BP. Clear, independent evidence that the severe climatic deterioration of Holocene Event 3 had an impact on the east Asia region at this time may be seen in marine sediment records from the South China Sea, the East China Sea and the North-west Pacific generally. Jian et al. (1996) used oxygen isotope analysis on planktonic foraminifera from deep water marine sediments in the East China Sea to reconstruct sea-surface

temperatures during the transition to the late Holocene, observing major cooling at the start of the Neoglacial around 4200 cal BP. Sun et al. (2005)'s oxygen isotope analyses from corals in the South China Sea show a major weakening of the monsoon and increase in its variability during the same period. Wang et al. (1999a, b), also using oxygen isotope data, identified a very clear abrupt cooling, the most severe in the Holocene apart from the 8200 cal BP event, at 4200-4000 cal BP in the northern South China Sea, a feature also observed by Wei et al. (1998) and by Chinzei et al. (1987) in the Sea of Japan. The ca. 4200 cal BP major cooling event is well attested in proxy climate records from the east Asia region, and it coincides with the apparent collapse of the late Neolithic cultures of the lower Yangtze area.

Although the circumstantial evidence that environmental pressures of varying kinds led to the collapse of the Liangzhu culture of the lower Yangtze around 4200 cal BP is very persuasive, we still do not know in detail the vegetation changes that preceded and accompanied the 4200 cal BP event there. New, high resolution palaeoecological data are required from within the Liangzhu's core settlement and agricultural area with which to establish these environmental preconditions. Most previous palynological research has been of low resolution, or situated too close to archaeological sites to be able to separate clearly any cultural impacts from the background vegetation history, or too far away to provide evidence of conditions within the cultural heartland itself. Also, not every pollen record agrees with the hypothesis that agricultural production almost ceased in the Taihu lowlands at this time (Itzstein-Davey et al., 2007b). In this paper we use high resolution palynology (both pollen and non-pollen palynomorphs) to investigate the nature and severity of the environmental changes during the transition to the Neoglacial in the Yangtze coastal lowlands of eastern China, as expressed in vegetation patterns and hydrology. A site at Pingwang has been selected where a more regional pollen signal might

be expected, minimizing the influence of local human land-use in this intensively settled area, so that natural factors rather than agricultural impacts will have been the main driving force behind the vegetation history and environmental change. Zong et al. (2011) have shown that direct inundation by marine transgression could not have been the reason for the abandonment of the Taihu lowlands by Neolithic people at this time, and this paper will explore in detail whether climate deterioration and its consequences was the environmental driving force, if one existed.

2. The study area and site

The study area chosen is the coastal lowland around the Taihu lake west of Shanghai (Fig.1). Between the valley of the Yangtze river and Hangzhou Bay, throughout the mid- and Late Holocene this flat plain, mostly less than 5 m above sea level, was naturally occupied by woodland and by wetland ecosystems which had developed during and after the main postglacial sea-level rise on this coast (Tao et al. 2006; Zong et al., 2011, 2012a, b; Wang et al., 2012). Higher ground to the west and north (Fig. 1a) and low hills within the wetland would have naturally carried forest. As well as the main Lake Tai (Qu, 2000; Wang et al., 2001) these wetlands included many other lakes and smaller water bodies and a wide expanse of peat-forming bogs, swamps and marshes. Although sheltered from direct marine inundation after the mid-Holocene by a system of coastal barrier ridges (Yan et al., 1989), these wetland systems remained affected by the groundwater influence of sea-level fluctuation (Zong et al., 2011). While their fertile marsh soils made them highly attractive for Neolithic settlement and intensive wet rice agriculture under the warm and wet climate of the mid-Holocene Megathermal phase (Cao et al., 2006; Lu, 2007; Atahan et al., 2008; Qin et al., 2011), the Taihu lowlands would have been highly vulnerable to any changes to their hydrology, sedimentation regime and natural

160 vegetation cover caused by even small-scale fluctuations in sea level or climate, or by fluvial
161 input from the nearby Yangtze river. Lying at the biogeographical boundary between
162 northeastern China's warm temperate broadleaf forests and southeastern China's sub-tropical
163 evergreen forests (Box, 1995; Li et al., 1995; Ren and Beug, 2002; Ren, 2007; Cao et al., 2013),
164 the natural vegetation of the lower Yangtze region would have been sensitive (Guiot et al., 2008)
165 to climatic deterioration caused by the proposed reduction of the strength of the EAM during the
166 transition to the Late Holocene (Morrill et al., 2003; Wang et al., 2005). It is an ideal location to
167 test the environmental consequences of climate change during the transition to the Late Holocene
168 Neoglacial in east China, and provide firmer evidence to assist interpretation of the
169 archaeological record and evaluate the putative cultural responses (Chen et al., 2005, 2008). It
170 can be difficult to distinguish vegetation changes caused by major climatic change from those set
171 in train by human land-use (Liu and Qiu, 1994), particularly if both may be occurring at the same
172 time and place, as both involve disturbance of established plant communities and the
173 regeneration of changed vegetation units through seral pathways. This is particularly so in the
174 Taihu lowlands, where usually population levels were high and agricultural activities were
175 intensive during the mid- to late Holocene (Wu et al., 2012). For the purposes of this study a site
176 was required that was at a significant distance from known Neolithic centres of agricultural
177 activity, identified by archaeological sites and their concentrations of cultural material that are
178 usually situated upon slightly higher land in the wetland. This should therefore contain a more
179 'natural' environmental signal with the influence of anthropogenic impacts on the vegetation
180 reduced as much as possible, although there would still be a regional cultural signature in the
181 pollen rain. It is not easy to get far away from known Neolithic cultural sites in this area, as there
182 are so many. Nevertheless such locations do exist, mainly in areas that were very wet and

perhaps agriculturally marginal, and one was selected at Pingwang to the south-east of Lake Taihu (Fig. 1) which lay more than 10 km from the nearest known archaeological site, and even further from any significant Neolithic settlement. The local hydrological history at Pingwang (30° 57' 30"N; 120° 38' 25"E; altitude 1.6 m Yellow Sea Datum) was summarised by Zong et al. (2011, 2012b). The sediments (Table 1) cover the whole of the mid- and late Holocene and record the development of various freshwater habitats above estuarine sediments after the postglacial rise of sea level was complete (Zheng and Li, 2000; Wang et al., 2012) and the final withdrawal of estuarine conditions from the area by 7000 cal BP after the establishment of coastal barrier ridges. Radiocarbon dating showed that the profile included the period of Liangzhu settlement and the Neoglacial cultural hiatus.

3. Material and methods

Samples were prepared for palynological analysis at 5 cm intervals using standard laboratory techniques, including alkali digestion, sieving at 180 µm, hydrofluoric acid digestion and acetolysis (Moore et al., 1991). Microfossils (all palynomorphs <180 µm in size) were identified using reference keys and type slides and counted using a stereomicroscope at magnification of ×400, using ×600 oil immersion lenses for critical features. Identification of pollen grains followed Wang et al. (1995) and pteridophyte spores Zhang et al. (1990). Where very similar pollen taxa cannot be differentiated with certainty, they are shown as a composite genus, i.e. *Ulmus/Zelkova*, *Corylus/Ostrya* and *Castanopsis/Lithocarpus*. *Quercus* includes *Lepidobalanus*. *Typha angustifolia* includes *Sparganium*. A minimum of 200 land pollen grains was counted at each sampled level, plus all aquatic pollen and pteridophyte and bryophyte spores observed while attaining that sum. Non-pollen palynomorphs (NPPs), mainly comprising fungal spores

and algae, were also recorded with at least 200 identified on the pollen slides at each level. Taxonomic identification of NPPs was achieved where possible, otherwise they were identified using the catalogue of Type (HdV) numbers at the Hugo de Vries laboratory, Amsterdam, using illustrations and descriptions published in several papers (e.g. van Geel, 1986, 2001; van Geel and Aptroot, 2006). Microscopic charcoal particles (microcharcoal) were counted upon the microscope slides relative to the pollen sum, providing a pollen/microcharcoal ratio.

Our dates on the Pingwang profile comprise AMS radiocarbon results and in this study, as in most other recent research in this area, only fragile terrestrial plant macrofossils, pollen residues or basal peat that has accumulated in situ have been dated, bulk alluvial sediment being avoided. Pollen residues have been shown in recent published studies from the area (Itzstein-Davey et al., 2007a; Atahan et al., 2008; Qin et al., 2011) to provide reliable age curves. A statistically significant inversion occurs low in the profile in Table 2, but these dates refer to the early Holocene Megathermal which is not the main focus of this study. It is considered that the pollen residue date at 321 cm depth is more likely to be correct, based on existing regional pollen and radiocarbon evidence. The upper dates, bracketing the age period of interest in this study, are consistent and are accepted as accurate. Dates were calibrated according to Calib6.1 (Reimer et al., 2009) using the IntCal09 programme. Microfossil diagrams were constructed using the TILIA program TGView (Grimm, 2004).

4. Results and interpretation

While Zong et al. (2012a) showed only selected summary curves from Pingwang, here we present full microfossil data. Tree and shrub pollen percentages are shown on Fig. 2, herb pollen and pteridophyte spores on Fig. 3 and NPPs on Fig. 4, all with frequencies calculated as

percentages of the total arboreal (tree + shrub) sum that reflects the more regional pollen rain. Fig. 2 includes summary curves for trees and shrubs based upon their life-form and temperature tolerances, although warm temperate and cool temperate are not separated. The CONISS (constrained incremental sum of squares) cluster analysis function (Grimm, 1987) within TGVView has been used to zone the diagram based on the total land pollen percentages, and has recognised four pollen assemblage zones (PW-a to d) which are applied to all of the diagrams. The CONISS dendrogram is shown on Fig. 2.

Zone PW-a (380-362.5 cm). This zone is dominated by the subtropical tree *Cyclobalanopsis* and the warm temperate trees *Quercus* and *Salix*, the latter perhaps abundant in local carr habitat with its locally sourced pollen frequencies depressing *Cyclobalanopsis* percentages. There are background levels of temperate taxa such as *Ulmus/Zelkova*, *Castanea* and *Liquidambar*. Cooler climate taxa are poorly represented, with only *Pinus* pollen significant, probably not locally sourced. Poaceae, Cyperaceae and *Typha angustifolia* dominate the herb assemblage. Prominent NPP types include Sordariaceae (55A), *Coniochaeta* cf. *ligniaria* and HdV-11, as well as marsh/reedswamp taxa HdV-306 and 708. This zone corresponds to the basal peat layer at the site (Table 1) and includes evidence of carr and shallow water marsh environments. Zong et al. (2011) recorded a mixed diatom assemblage in this lower part of the profile, with evidence of saltmarsh or estuarine influence. The non-wetland vegetation was a subtropical forest with temperate elements which the date suggests existed not long after the start of the Megathermal mid-Holocene climate optimum.

Zone PW-b (362.5-232.5 cm). This zone is dominated by the subtropical genera *Cyclobalanopsis* and *Castanopsis/Lithocarpus* which consistently provide 50% of arboreal pollen, with temperate deciduous *Quercus* accounting for most of the rest. Other temperate trees

are also significant, including *Ulmus/Zelkova*, *Castanea*, *Pterocarya* and *Liquidambar*. Conifers and cold tolerant trees are poorly represented, with a low background curve for *Pinus*. Non-arboreal pollen and spores are low, supplied mainly by Poaceae and Cyperaceae. Aquatic taxa are important, and *Potamogeton* and *Ceratopteris* increase later in the zone and there is a peak of the aquatic *Salvinia*. NPPs become increasingly dominated by aquatic types, with shallow water marsh taxa like *Zygnema*, *Mougeotia*, HdV-306 and HdV-708 being replaced in the upper part of the zone by more open water algal types, especially *Pediastrum*, HdV-128 and the cyanobacterium *Gloeotrichia*. The regional forest comprised a typical Megathermal optimum assemblage with evergreen subtropical trees dominant but with a significant temperate component. The local environment changed from shallow water marsh to aquatic and eutrophic, biologically productive habitats with deeper standing water, as shown by *Pediastrum* abundance, nearer to 5000 cal BP. Such *Pediastrum* frequencies are typical of limnic environments with emergent aquatic vegetation under warm, mid-Holocene climates (Jankovská and Komárek, 2000).

Zone PW-c (232.5-177.5 cm). At the start of this zone subtropical tree frequencies, *Cyclobalanopsis* and *Castanopsis/Lithocarpus*, fall from 50% to 25% of arboreal pollen. Of the warm temperate trees, the more cool-tolerant *Betula*, *Fagus* and *Alnus* increase but the more thermophilous, including *Pterocarya* and *Castanea*, decline markedly. Against this trend pollen of Moraceae, usually subtropical trees in east China (Sun et al., 2003), appears and rises in frequency in mid-zone. Distinct increases occur in the cold-tolerant trees Cupressaceae (which includes *Juniperus*), *Picea* and *Pinus*. Frequencies of the main warm temperate forest trees *Quercus*, *Liquidambar* and *Ulmus/Zelkova* are unaffected. The summary curves on Fig.1 show an almost direct replacement of the subtropical genera by the cold-tolerant taxa. There is little

change in the non-arboreal pollen and spores except for peaks in *Potamogeton*, *Myriophyllum* and Adiantaceae, all floating aquatics. The NPP assemblage shows major changes with *Gloeotrichia* rising to dominance and *Pediastrum* reduced, although still important. Peaks of the cyanobacterium *Anabaena*, intolerant of high temperatures, accompany *Gloeotrichia*'s rise. The local aquatic habitat seems to have become colder, deeper and less eutrophic, while the regional forest also adapted to colder temperatures during this zone.

Zone PW-d (177.5-130 cm). At the start of this zone, which has an interpolated date of 4200 cal BP, the frequencies for the subtropical trees *Cyclobalanopsis* and *Castanopsis/Lithocarpus* continue to fall, dropping to 15% of arboreal pollen. Moraceae, however, maintains values of almost 10% of total tree and shrub pollen. Warm temperate trees are little changed, but the cold-tolerant and coniferous taxa rise to 35% of the total. *Pinus* and Cupressaceae are particularly increased, and Taxodiaceae is consistently recorded. Aquatic pollen is still important although the rise of *Typha* and *Ceratopteris* late in the zone implies falling water levels. Present in low frequencies throughout, *Artemisia* increases late in the zone. *Gloeotrichia* is still abundant early in the zone, but decreases as marsh herbs, ferns and NPPs increase after ca 2800 cal BP. This zone sees a continued expansion of conifers in the forest at the expense of the subtropical genera.

5. Discussion

5.1. Yangtze region vegetation history

The biogeographical location of the lower Yangtze region means that its natural vegetation contains elements of both subtropical and temperate forests, and during the mid-Holocene

Megathermal optimum phase, subtropical forest trees were common there as the monsoon front lay to the north and the Yangtze had a very warm, humid and wet climate (Shi et al., 1993; Zheng et al., 2004; Zong et al., 2007; Innes et al., 2009; Li et al., 2010a; Cao et al., 2013). *Castanopsis* and *Cyclobalanopsis* were the dominant taxa, associated with thermophilous deciduous trees *Quercus*, *Liquidambar* and *Ulmus/Zelkova*. There are several published pollen profiles from the lower Yangtze region with which the Pingwang Megathermal record can be compared (Liu et al., 1992; Xu et al., 1996; Chen et al., 2005; Tao et al., 2006; Yi et al., 2006; Shu et al., 2007; Innes et al., 2009; Li et al., 2010b) as well as more general regional syntheses for east China (Liu, 1988; Sun and Chen, 1991; Ren and Beug, 2002; Zhang et al., 2005; Ren, 2007), and all show this general forest history. Zhao et al. (2009) have analysed the data from many pollen records across all of EAM China and conclude that whatever the Megathermal forest type, there was a shift to a more cold-tolerant forest community, in many regions with major expansion of conifers in particular, after about 4200 cal BP, and it may well have been responsible for the major forest decline that occurred in more arid regions (Zhao et al., 2009; Herzsuh et al., 2010), instead of the human impact explanation favoured by earlier workers (Liu and Qiu, 1994; Ren, 2000). This climate shift is well represented in the lower Yangtze (Wu et al., 2012), with a broadleaf temperate and conifer woodland established. As most rice agriculture occurred here on sites reclaimed from wetlands (Li et al., 2012), human impacts on the forest were probably not great except close to settlement sites (Atahan et al., 2008). This remained the case until major human forest clearance occurred after ca 2500 cal BP. (Atahan et al., 2007; Wang et al., 2011).

5.2. The 4200 cal BP climatic deterioration in China

319 While reservations exist (Maher, 2008; Maher and Thompson, 2012), the severe climatic
320 deterioration at ca 4200 cal BP appears to be well attested in speleothem records from caves in
321 several areas of China (Wang et al., 2001; Dykoski et al., 2005; Shao et al., 2006; Cosford et al.,
322 2008; Hu et al., 2008; Dong et al., 2010). This widespread switch to cold conditions (Shi et al.,
323 1993) is substantiated by many other forms of proxy data (Hong et al., 2000; Li et al., 2010b),
324 including pollen records (Xiao et al., 2004; Lim and Fujiki, 2011; Chen et al., 2012). In more
325 northerly and westerly areas of China towards the EAM margins the 4200 cal BP climatic
326 deterioration and instability was expressed by a rapid switch to much more arid conditions (Hong
327 et al., 2001; Schettler et al., 2006; Jiang and Liu, 2007; Mischke and Zhang, 2010; Wang et al.,
328 2010; Wen et al., 2010), as the summer monsoon weakened and the winter monsoon
329 strengthened (Yu et al., 2006; Jiang and Liu, 2007; Yancheva et al., 2007; Yang and Scuderi,
330 2010). Precipitation would have been the dominant factor controlling vegetation distributions
331 (Wen et al., 2013). In the western part of the Chinese Loess Plateau, the climate until 4200 cal
332 BP was humid (An et al., 2003; 2004; Gao et al., 2007), with organic palaeosol development.
333 Southeast of Lanzhou, for example, wetlands seem to have existed in several river valleys,
334 adjacent to Neolithic settlements (Feng et al., 2004, 2006). At about 4200 cal yr BP, the climate
335 suddenly became drier in this region and from the reduction in the number of archaeological sites
336 from this time, An et al. (2004, 2005) concluded that this period of intense aridity, which lasted
337 for some centuries, had a major effect on the Neolithic Qijia people who lived in these river
338 valleys, causing the collapse of their farming culture (Mo et al., 1996; Liu et al., 2010a).
339 Previously distinct pollen and charcoal records of major human land-use impacts in this area also
340 terminate at this time (Li et al., 2012). At the western and northern margins of the area of
341 monsoonal influence, this period of maximum aridity (Wei and Gasse, 1999) caused lakes to

dessicate almost completely (Chen et al., 1991; Morrill et al., 2006; Zhai et al., 2011). This drying tendency since about 4900 cal BP extended even to eastern coastal China, where some lake levels started to fall (Wu et al., 2010).

One of the clearest expressions of the climatic deterioration is in vegetation records (Zhao et al., 2009), many of which show major changes in the centuries leading up to 4000 cal BP. On the Western Loess Plateau and in the uplands of central China there was an abrupt change from forest to steppe as a response to the greatly increased aridity (Feng et al., 2006; Zou et al., 2009; Herzsuh et al., 2010; Zhou and Li., 2012). Elsewhere in north and west China at this time this much colder and arid phase changed mixed forest to domination by conifers (Sun and Weng, 1992; Jarvis, 1993; Ren, 2000; Makohonienko et al. 2004; Xiao et al., 2004; Jiang and Liu, 2007; Liu et al., 2010b; Xu et al., 2010; An et al., 2012), as pine was particularly favoured over oaks under the more arid and cold conditions (Ren and Zhang, 1998; Yi et al., 2003). Although there has been discussion as to whether human activity or climate change was responsible for the spread of *Pinus* in northeast China around 4200 cal BP (Ren, 2000; Xu et al., 2010), there is little evidence for human interference with the woodland during that period and climate change seems to be the most likely cause. In central eastern China the evergreen trees common during the Megathermal (Shi et al., 1994) were everywhere displaced by conifers within mixed, cool temperate woodland (Ren and Beug, 2002; Zhu et al., 2010), a change to a cooler forest biome recorded even in south China and Taiwan (Liew et al., 2006; Wang et al., 2007; Lee et al., 2010; Wu et al., 2012). There is more persuasive evidence for human activity having had an influence on the forest in the Yangtze region (Chen et al., 2009; Li et al., 2010b; Wu et al., 2012), but it declines and then stops almost everywhere in the centuries leading up to the 4200 cal BP event.

In contrast to the aridity phase in north and west China, in many places at this time in eastern and southern China peat deposits begin to form or expand markedly (Zhao et al., 2007; Zhang et al., 2011), including in the Taihu lowlands (Zhu et al., 2006), under greatly wetter conditions, providing pollen and other proxy records that confirm the abrupt and major climatic cooling at ca 4200 BP (Hong et al., 2000; Yu et al., 2006; Zhao et al., 2007; Ma et al. 2008, 2009) following the southerly withdrawal and weakening of the EAM (Wang et al., 2005; Liu et al., 2010c) which coincided with stronger winter monsoons (Yancheva et al., 2007). An important consequence of the 4200 cal BP climatic decline in eastern China seems to have been a dramatic increase in the incidence and severity of flooding in river valleys (Tan et al., 2008; Wu et al. 2012), where much of Late Neolithic settlement and agriculture was concentrated. Fields and settlements were often located close to water sources, lakes and river channels, making them highly vulnerable to rapid fluctuations in water levels (Wu et al., 2010). Often cultural deposits are terminated by major flood sediment layers that date to this event (Huang and Zhang, 2000; Bai et al., 2008; Yao et al., 2008; Huang et al., 2010, 2011, 2012a, b; Zhang et al., 2010) and represent disasters that must have buried fields and forced settlement evacuation (Xia et al., 2003; Liu et al., 2012), including some in the Yangtze Delta itself (Zhang, 2007; Zhang et al., 2004a) and at the termination of the Liangzhu (Zhu et al., 1996; Zhang et al. 2004b; Shi et al., 2008). Huang et al. (2011) record repeated floods which must have made cultural recovery in these locations very difficult.

5.3. *Environmental evidence at Pingwang*

5.3.1. *Hydrological evidence*

386 After the withdrawal of brackish marsh conditions from Pingwang, the site became dominated
387 throughout its history by freshwater wetland environments, and increasingly by fully aquatic
388 conditions, which probably explains the absence of archaeological material nearby and the lack
389 of palynological evidence for agriculture throughout the profile (Figs. 3 and 4). The NPP record
390 is particularly informative regarding the local hydrology, with HdV-128, *Spirogyra*,
391 *Gloeotrichia*, *Zygnema* and *Pediastrum* all abundant and reflecting open water of various depths
392 and trophic status, and the aquatic herb taxa *Potamogeton*, *Typha* and *Salvinia* support this. The
393 significance of HdV-128, *Zygnema* and HdV-708 in zone PW-b suggests mesotrophic open
394 water of relatively shallow depth (Bakker & van Smeerdijk, 1982; Pals et al., 1980). The
395 microcharcoal curve is not high and probably represents a background regional signal of human
396 activity, presumably on drier land but perhaps also burning of carr scrub within the wetland to
397 establish paddy fields (Innes et al. 2009) and of dead plant stubble material to maintain them
398 (Cao et al. 2006; Dodson et al., 2006; Zheng et al., 2009; Li et al., 2012; Hu et al., 2013; Zhuang
399 et al., 2014). Although moderate throughout, the increased values at the top and base of the
400 profile correlate with periods of shallower-water marsh conditions, when some increase in local
401 human presence might be expected. It is interesting that microcharcoal frequencies fall sharply
402 during the episode at the end of zone PW-c correlated with the 4200 cal BP event, supporting
403 theories of greatly reduced human activity throughout the Taihu plain. The non-arboreal pollen
404 and the NPPs indicate deeper water at this time, with *Gloeotrichia* abundant and replacing
405 *Pediastrum* as a deeper and cooler but perhaps more eutrophic (van Geel et al., 1996) aquatic
406 environment developed. Some eutrophication due to drainage from agricultural land during the
407 Liangzhu period in zone PW-c might be expected, and low peaks of *Anabaena* would support
408 this (van Geel et al., 1994; Hillbrand et al., 2014), although the major bloom in *Gloeotrichia*

would suppress the abundance of the light-demanding *Anabaena*. *Gloeotrichia* might also have been favoured by rising water levels and thus clearer water at this time, being favoured by increased light levels (Chmura et al., 2006). Presumably Pingwang was subsumed within the expanded water bodies of Taihu and the other lakes in the area which came into being after 4600 cal BP (Wang et al., 2001; Zong et al., 2012a, b), following significantly increased rainfall and drainage discharge from the Yangtze valley (Long et al., 2014; Wu et al., 2014).

5.3.2. Climatic or human influence

The arboreal pollen curves at Pingwang record woodland history all through the mid-Holocene Megathermal, across the Late Holocene transition and into recent times, and providing evidence from beyond the local aquatic environments at Pingwang. The expansion of conifers, mainly *Pinus*, and the decline of subtropical evergreens after ca 5400 cal BP is very clear and is the important feature of the vegetation history. In this respect it records the gradual onset of the Neoglacial in eastern China at this time as recorded in other sites (Zhong et al., 2010). The choice of the Pingwang site was designed to reduce the human element in woodland history as far as possible, as there are no records of significant Neolithic settlement near the site, even at the time of the greatest Liangzhu expansion. A complete absence of human influence on the site cannot be assumed, however, as there would be few areas beyond all human activity in the intensively settled Taihu lowland, if any, even the constantly flooded Pingwang locality. The complete absence of any big grass pollen grains that could be attributed to rice in the Pingwang pollen record, however, suggests that the site was well away from any cultivation throughout the Neolithic period. There are also no NPP indicators of pastoral farming and concentrations of

432 livestock such as the dung fungi *Sporormiella* or *Podospora*, as were found near the early
433 Neolithic agricultural settlement at Kuahuqiao (Zong et al., 2007; Innes et al., 2009). All are
434 absent from Pingwang, and virtually all NPPs reflect wetland environments. It is possible that the
435 rise in *Pinus* and Moraceae frequencies during zone PW-c might have been caused by
436 regeneration of secondary woodland after forest clearance in this lower Yangtze region (Huang
437 and Zhang, 2000; Chen et al., 2009; Li et al., 2010b), as this is the main Liangzhu settlement and
438 farming period and area, and some authors have reported extensive forest clearance and
439 agricultural intensification in the Late Neolithic in the lower Yangtze catchment between 5000
440 and 4200 cal BP, when it abruptly stopped (Yasuda et al., 2004; Zhuang et al., 2014). Liu (1988),
441 Liu et al. (1992), Liu and Qiu (1994) and Okuda et al. (2003) have suggested that both climatic
442 cooling and human disturbance may have contributed to the late Holocene *Pinus* rise in the lower
443 Yangtze. This cannot be the case at Pingwang, however, where effects of significant forest
444 clearance would be visible in the pollen record, even if none took place near to the site. During
445 zone PW-c, when the Liangzhu culture was flourishing, tree pollen values generally remain high,
446 with no increase in non-tree pollen that might indicate deforestation. The rise of Moraceae in
447 zone PW-c is strange, however, if subtropical tree genera were in progressive decline. It is
448 possible that Moraceae genera, mainly secondary trees, were increasing locally in successional
449 woodlands as populations of *Cyclobalanopsis* and *Castanopsis/Lithocarpus* declined. An
450 alternative, however, is that the Moraceae curve reflects human activity of a particular kind,
451 taking advantage of the extensive water body at Pingwang. Retting of cannabis for fibres (e.g.
452 Schofield and Waller, 2005) in the shallow lake edges could account for the increase in
453 Moraceae pollen when subtropical genera are otherwise in decline. Moraceae frequencies are
454 also high in zone PW-d, in later prehistory, when human activity might also be expected,

although there remains no sign of rice agriculture. Similar high Moraceae frequencies attributed to human activity have been recorded elsewhere in the Taihu region (Itzstein-Davey et al., 2007b). It is interesting that Moraceae percentages fall abruptly at the end of zone PW-c, during the end-Neolithic cold phase and cultural hiatus, before recovering later. This would support both the origin of the Moraceae pollen in a human activity and the effectiveness of the ca. 4200 cal BP cold phase in temporarily stopping it, along with all other human activity and settlement in the Taihu plain.

The percentage representation of temperate trees, including the forest co-dominant *Quercus*, is almost unchanged throughout the Pingwang pollen profile, and forest composition changes are caused dominantly by the replacement of sub-tropical genera by cold-tolerant trees, mainly conifers. It is unreasonable to suggest that human land clearance for farming would have affected only sub-tropical trees and leave unscathed the deciduous temperate trees, especially the abundant oaks, that grew alongside them in the mixed forest. Only climate change and colder conditions can account for the direct replacement of the sub-tropical component of the forest by pine and other conifers. Although closer to the lower end of their temperature tolerance ranges, the deciduous temperate trees like *Quercus* could withstand the new colder environment and survive, whereas the sub-tropical genera could not. Given a competitive advantage by the colder climate, conifers moved in to occupy the place in the forest community vacated by the subtropical evergreens, particularly after the abrupt shift to the Neoglacial at ca 4200 cal BP, while the proportion of temperate deciduous trees remained unchanged.

6. Conclusions

476 The profile from Pingwang in the centre of the Taihu coastal plain contains no pollen evidence of
477 cultivation, other forms of human land-use or clearance of woodland, except perhaps for the
478 equivocal evidence of a slight increase in open ground weeds after ca 2500 cal BP. Certainly
479 there is no indication of agricultural activity there throughout the Late Neolithic, despite the
480 intensive settlement and agricultural use of the Taihu lowlands by the Liangzhu culture. Because
481 of its palaeogeography, Pingwang therefore provides a record of sub-regional forest history,
482 uncomplicated by local human activity, unless the Moraceae pollen curve indicates fibre
483 processing in the lake. It shows very clearly the rise of *Pinus* and other conifers after ca 5400 cal
484 BP, and the rapid acceleration of that rise at ca 4200 cal BP. These two dates agree very well
485 with the timings of the onset of climatic deterioration and its severe intensification noted around
486 the globe (e.g. Geirsdóttir et al., 2013). It also clarifies the reasons for that radical transformation
487 of the forests of the lower Yangtze region, and by extension for the same changes in forest
488 composition that can be seen throughout eastern China at this time, with change beginning at ca
489 5400 cal BP. Although it has been suggested previously that the rise of pine and other
490 successional tree taxa in the Late Neolithic might have resulted from regeneration after major
491 deforestation for agriculture, the Pingwang data make clear that the gradual and then sudden
492 switch to a cold Neoglacial climate can account for the forest changes observed in regional
493 vegetation history. The increased dominance of fully aquatic conditions across the Late
494 Holocene transition at Pingwang is shown particularly clearly by the NPP results and accords
495 with the evidence from several other Yangtze valley sites for severe flooding events at this time.
496 It supports the hypothesis that severe and regular freshwater flooding or rising local water level
497 (Stanley et al., 1999; Zong et al., 2012a; Long et al., 2014) on the Taihu plain, allied to
498 significantly lower temperatures, is the explanation for the collapse of the Liangzhu culture, as

well as other Late Neolithic cultures of the Yangtze valley, at ca 4200 cal BP. The abrupt switch to such adverse conditions would have made organised rice farming and complex settlement virtually impossible until a measure of climatic amelioration occurred some centuries later.

Acknowledgements

This research was funded by the Leverhulme Trust, U.K (project number F/000128/AL). We are grateful to Dr. J. Chen of the Shanghai Museum for his advice regarding the location of Neolithic sites in the study area. Chris Orton of the Design and Imaging Unit, Geography Department, Durham University produced the figures.

References

An, C., Feng, Z., Tang, L., 2003. Evidence of a humid mid-Holocene in the western part of the Chinese Loess Plateau. *Chin. Sci. Bull.* 48, 2472-2479.

An, C.B., Feng, Z., Tang, L., 2004. Environmental change and cultural response between 8000 and 4000 cal. Yr B.P. in the western Loess Plateau, northwest China. *J. Quat. Sci.* 19, 529-535.

An, C-B., Tang, L., Barton, L., Chen, F-H., 2005. Climate change and cultural response around 4000 cal yr B.P. in the western part of Chinese Loess Plateau. *Quat. Res.* 63, 347-352.

An, C-B., Lu, Y., Zhao, J., Tao, S., Dong, W., Li, H., Jin, M., Wang, Z., 2012. A high-resolution record of Holocene environmental and climatic changes from Lake Balikun (Xinjiang, China): implications for central Asia. *Holocene* 22, 45-52.

521

522 An, Z.S., 2000. The history and variability of the east Asian paleomonsoon climate. *Quat. Sci.*

523 *Rev.* 19, 171-187.

524

525 An, Z.S., Porter, S.C., Kutzbach, J.E., Wu, X., Wang, S., Liu, X., Li, X., Zhou, W., 2000.

526 Asynchronous Holocene optimum of the East Asian monsoon. *Quat. Sci. Rev.* 19, 743-762.

527

528 Anderson, A., Gagan, M., Shulmeister, J. 2007. Mid-Holocene cultural dynamics and climatic

529 change in the Western Pacific. In: Anderson, D.G., Maasch, K.A., Sandweiss, D.H. (Eds.)

530 *Climate change and Cultural Dynamics: a Global Perspective on mid-Holocene Transitions.*

531 Academic Press, London, pp. 265-296.

532

533 Atahan, P., Grice, K., Dodson, J. 2007. Agricultural and environmental change at Qingpu,

534 Yangtze Delta region, China: a biomarker, stable isotope and palynological approach. *Holocene*

535 17, 507-515.

536

537 Atahan, P., Itzstein-Davey, F., Taylor, D., Dodson, J., Qin, J., Zheng, H., Brooks, A., 2008.

538 Holocene-aged sedimentary records of environmental changes and early agriculture in the lower

539 Yangtze, China. *Quat. Sci. Rev.* 27, 556-570.

540

541 Bai, J., Zhou, H., Zhu, C. 2008. Archaeological discovery and research on the remains of the

542 palaeoflood of the Yuxi site. *Chin. Sci. Bull.* 53 (Supp. 1), 18-27.

543

544 Bakker, M., van Smeerdijk, D.G., 1982. A palaeoecological study of a late Holocene section
545 from “Het Ilperveld”, western Netherlands. *Rev. Palaeobot. Palynol.* 36, 95-163.
546

547 Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M.N., Showers, W., Hoffmann, S., Lotti-
548 Bond, R., Hajdas, I., Bonani, G., 2001. Persistent solar influence on north Atlantic climate during
549 the Holocene. *Science* 294, 2130–2136.
550

551 Booth, R., Jackson, S.T., Forman, S.L., Kutzbach, J.E., Betis III, E.A., Kreigs, J., Wright, D.K.,
552 2005. A severe centennial-scale drought in mid-continental North America 4200 years ago and
553 apparent global linkages. *Holocene* 15, 321–28.
554

555 Box, E.O., 1995. Climate relations of the forests of east and southeast Asia. In: Box E.O, et al.
556 (Eds.), *Vegetation Science in Forestry*. Kluwer Academic Publishers, Dordrecht, pp23-55.
557

558 Cai, Y., Tan, L., Cheng, H., An, Z., Edwards, R.L., Kelly, M.J., Kong, X., Wang, X., 2010. The
559 variation of summer monsoon precipitation in central China since the last deglaciation. *Earth and*
560 *Planetary Science Letters* 291, 21–31.
561

562 Cao, X., Ni, J., Herzsuh, U., Wang, Y., Zhao, Y., 2013. A late Quaternary pollen dataset from
563 eastern continental Asia for vegetation and climate reconstructions: set up and evaluation. *Rev.*
564 *Palaeobot. Palynol.* 194, 21-37.
565

566 Cao, Z., Ding, J., Hu, Z., Knicker, H., Kögel-Knabner, I., Yang, L., Yin, R., Lin, X., Dong, Y.,
567 2006. Ancient paddy soils from the Neolithic age in China's Yangtze River Delta.
568 *Naturwissenschaften* 93, 232–236.

569

570 Caseldine, C.J., Turney, C., 2010. The bigger picture: towards integrating palaeoclimate and
571 environmental data with a history of societal change. *J. Quat. Sci.* 25, 88-93.

572

573 Chen, F.H., Shi, Q., Wang, J.M., 1999. Environmental changes documented by sedimentation in
574 Lake Yiema in arid China since the late glaciation. *J. Paleolimnol.* 22, 159–169.

575

576 Chen, W., Wang, W-M., 2012. Middle-Late Holocene vegetation history and environment
577 changes revealed by pollen analysis of a core at Qingdao of Shandong Province, East China.
578 *Quat. Int.* 254, 68-72.

579

580 Chen, W., Wang, W., Dai, X., 2009. Holocene vegetation history with implications of human
581 impact in the Lake Chaohu area, Anhui Province, East China. *Veget. Hist. Archaeobot.* 18, 137-
582 146.

583

584 Chen, Z., Wang, Z., Schneiderman, J., Tao, J., Cai, Y., 2005. Holocene climate fluctuations in
585 the Yangtze delta of eastern China and the Neolithic response. *Holocene* 15, 915-924.

586

587 Chen, Z., Zong, Y., Wang, Z., Wang, H., Chen, J., 2008. Migration patterns of Neolithic
588 settlements on the abandoned Yellow and Yangtze River deltas of China. *Quat. Res.* 70, 301-
589 314.

590

591 Chinzei, K., Koike, H., Oba, T., Matsushima, Y., Kitazato, H., 1987. Secular changes in the
592 oxygen isotope ratios of mollusc shells during the Holocene of central Japan. *Palaeogeogr.*
593 *Palaeoclimatol. Palaeoecol.* 61, 155-166.

594

595 Chmura, G., Stone, P.A., Ross, M.S., 2006. Non-pollen microfossils in Everglades sediments.
596 *Rev. Palaeobot. Palynol.* 141, 103-119.

597

598 Coombes, P., Barber, K., 2005. Environmental determinism in Holocene research: causality or
599 coincidence? *Area* 37, 303-311.

600

601 Cosford, J., Qing, H., Eglington, B., Matthey, D., Yuan, D., Zhang, M., Cheng, H., 2008. East
602 Asian monsoon variability since the Mid-Holocene recorded in a high-resolution, absolute-dated
603 aragonite speleothem from eastern China. *Earth Planet. Sci. Lett.* 275, 296-307.

604

605 Cullen, H.M., deMenocal, P.B., Hemming, S., Hemming, G., Brown, F.H., Guilderson, T.,
606 Sirocko, F., 2000. Climate change and the collapse of the Akkadian empire: Evidence from the
607 deep sea. *Geology* 28, 379–382.

608

Dalfes, H., Kukla, G., Weiss, H., Eds 1997. Third Millennium BC Climate Change and Old World Collapse, NATO ASI Series 1, vol. 49, Springer-Verlag, Berlin.

deMenocal, P.,B., 2001. Cultural responses to climate change during the Late Holocene. *Science* 292, 667-673.

Dodson, J.R., Hickson, S., Khoo, R., Li, X., Toia, J., Zhou, W., 2006. Vegetation and environment history for the past 14 000 yr BP from Dingnan, Jiangxi province, South China. *J. Integr. Plant Biol.* 48, 1018-1027.

Dong, J., Wang, Y., Cheng, H., Hardt, B., Eswards, R.L., Kong, X., Wu, J., Chen, S., Liu, D., Jiang, X., Zhao, K., 2010. A high resolution stalagmite record of the Holocene East Asian monsoon from Mt Shennongjia, central China. *Holocene* 20, 257-264.

Drysdale, R., Zanchetta, G., Hellstrom, J., Maas, R., Fallick, A., Pickett, M., Cartwright, I., Piccini, L., 2006. Late Holocene drought responsible for the collapse of Old World civilizations is recorded in an Italian cave flowstone. *Geology* 34, 101–104.

Dykoski, C.A., Edwards, R.L., Cheng, H., Yuan, D., Cai, Y., Zhang, M., Lin, Y., Qing, J., An, Z., Revenaugh, J., 2005. A high-resolution, absolute-dated Holocene and deglacial Asian monsoon record from Dongge Cave, China. *Earth Planet. Sci. Lett.* 233, 71–86

631 Feng, Z., Tang, L.Y., Wang, H.B., Ma, Y.Z., Liu, K-B., 2006. Holocene vegetation variations
632 and the associated environmental changes in the western part of the Chinese Loess Plateau.
633 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 241, 440-456.
634

635 Feng, Z., An, C.B., Wang, H.B., 2006. Holocene climatic and environmental changes in the arid
636 and semi-arid areas of China: a review. *Holocene* 16, 119-130.
637

638 Gao, H., Zhu, C., Xu, W., 2007. Environmental change and cultural response around 4200 cal.yr
639 BP in the Yishu River Basin, Shandong. *J. Geogr. Sci.* 17, 285–292.
640

641 Gear, A.J., Huntley B., 1991. Rapid changes in the range limits of Scots Pine 4000 years ago.
642 *Science* 251, 544-547.
643

644 Geirsdóttir, A., Miller, G., Larsen, D.J., Ólafsdóttir, S., 2013. Abrupt Holocene climate
645 transitions in the northern North Atlantic region recorded by synchronized lacustrine records in
646 Iceland. *Quat. Sci. Rev.* 70, 48-62.
647

648 Grimm, E.C., 1987. CONISS: a FORTRAN 77 program for stratigraphically constrained cluster
649 analysis by the method of incremental sum of squares. *Comp. Geosci.* 13: 13–35.
650

651 Grimm, E.C. 2004. TGView v. 2.0.2, Software. Illinois State Museum, Research and Collections
652 Center: Springfield, IL.
653

654 Guiot, J., Wu, H.B., Jiang, W.Y., Luo, Y.L., 2008. East Asian Monsoon and paleoclimatic data
 655 analysis: a vegetation point of view. *Clim. Past* 4, 137-145.
 656
 657 Gupta, A., Anderson, D., Overpeck, J., 2003. Abrupt changes in the Asian southwest monsoon
 658 during the Holocene and their links to the North Atlantic. *Nature* 421, 354–357.
 659
 660 He, Y., Theakstone, W.H., Zhang, Z., Zhang, D., Yao, T., Chen, T., Shen, Y., Pang, H., 2004.
 661 Asynchronous Holocene climatic change across China. *Quat. Res.* 61, 52-63.
 662
 663 Herzschuh, U., Birks, H.J.B., Liu X, Kubatzki, C., Lohmann, G. 2010. What caused the mid-
 664 Holocene forest decline on the eastern Tibet-Qinhai Plateau. *Glob. Ecol. Biogeogr.* 19, 278-286.
 665
 666 Hillbrand, M., van Geel, B., Hasenfratz, A., Hadorn, P., Haas, J.N., 2014. Non-pollen
 667 palynomorphs show human- and livestock-induced eutrophication of Lake Nussbaumersee
 668 (Thurgau, Switzerland) since Neolithic times (3840 BC). *Holocene* 24, 559-568.
 669
 670 Hong, Y.T., Jiang, H.B., Liu, T.S., Zhou, L.P., Beer, J., Li, H.D., Leng, X.T., Hong, B., Qin,
 671 X.G. 2000. Response of climate to solar forcing recorded in a 6000-year $\delta^{13}\text{O}$ time-series of
 672 Chinese peat cellulose. *Holocene* 10, 1–7.
 673
 674 Hong, Y.T., Wang, Z.G., Jiang, H.B., Lin, Q.H., Hong, B., Zhu, Y.X., Wang, Y., Xu, L.S., leng,
 675 X.T., Li, H.D., 2001. A 6000-year record of changes in drought and precipitation in northeastern

676 China based on a $\delta^{13}\text{C}$ time series from peat cellulose. *Earth and Planetary Science Letters* 185,
677 111–119.

678

679 Hong, Y.T., Hong, B., Lin Q.H., Shibata, Y., Hirota, M., Zhu, Y.X., Leng, X.T., Wang, Y.,
680 Wang, H., Yi, L., 2005. Inverse phase oscillations between the East Asian and Indian Ocean
681 summer monsoons during the last 12 000 years and paleo-El Niño. *Earth and Planetary Science*
682 *Letters* 231, 337-346.

683

684 Hu, C., Henderson, G.M., Huang, J., Xie, S., Sun, Y., Johnson, K.R., 2008. Quantification of
685 Holocene Asian monsoon rainfall from spatially separated cave records. *Earth and Planetary*
686 *Science Letters* 266, 221–32.

687

688 Hu, L., Chao, Z., Gu, M., Li, F., Chen, L., Liu, B., Li, X., Huang, Z., Li, Y., Xing, B., Dai, J.,
689 2013. Evidence for a Neolithic Age fire-irrigation paddy cultivation system in the lower Yangtze
690 River Delta, China. *J. Archaeol. Sci.* 40, 72-78.

691

692 Huang, C.C., Pang, J., Zha, X.C., Zhou, Y., Su, H., Li, Y., 2010. Extraordinary floods of 4100–
693 4000 a BP recorded at the Late Neolithic ruins in the Jinghe River Gorges, middle reach of the
694 Yellow River, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 289, 1-9.

695

696 Huang, C.C., Pang, J., Zha, X., Su, H., Jia, Y., 2011. Extraordinary floods related to the climatic
697 event at 4200 a BP on the Qishuihe River, middle reaches of the Yellow River, China. *Quat. Sci.*
698 *Rev.* 30, 460-468.

699

700 Huang, C.C., Pang, J., Zha, X., Zhou, Y., Su, H., Zhang, Y., Wang, H., Gu, H., 2012a. Holocene
701 palaeoflood events recorded by slackwater deposits along the lower Jinghe River valley, middle
702 Yellow River basin, China. *J. Quat. Sci.* 27, 485-493.

703

704 Huang, C.C., Pang, J., Zha, X., Zhou, Y., Su, H., Wan, H., Ge, B., 2012b. Sedimentary records
705 of extraordinary floods at the ending of the mid-Holocene climatic optimum along the Upper
706 Weihe river, China. *Holocene* 22, 675-686.

707

708 Huang, F., Zhang, M., 2000. Pollen and phytolith evidence for rice cultivation during the
709 Neolithic at Longqiuzhuang, eastern Jianghuai, China. *Veget. Hist. Archaeobot.* 9, 161-168.

710

711 Innes, J.B., Zong, Y., Chen, Z., Chen, C., Wang, Z. & Wang, H., 2009 Environmental history,
712 palaeoecology and human activity at the early Neolithic forager/cultivator site at Kuahuqiao,
713 Hangzhou, eastern China. *Quat. Sci. Rev.* 28, 2277-2294.

714

715 Itzstein-Davey, F., Atahan, P., Dodson, J., Taylor, D., Zheng, H., 2007a. Environmental and
716 cultural changes during the terminal Neolithic, Qingpu, Yangtze delta, eastern China. *Holocene*
717 17, 875-887.

718

719 Itzstein-Davey, F., Atahan, P., Dodson, J., Taylor, D., Zheng, H., 2007b. A sediment-based
720 record of Lateglacial and Holocene environmental changes from Guangfulin, Yangtze delta,
721 eastern China. *Holocene* 17, 1221-1231.

722

723 Jankovská, V., Komárek, J., 2000. Indicative value of *Pediastrum* and other coccal green algae
 724 in palaeoecology. Folia Geobot. 35, 59-73, 75-82.

725

726 Jarvis, D.I., 1993. Pollen evidence of changing Holocene monsoon climate in Sichuan Province,
 727 China. Quat. Res. 39, 325-337.

728

729 Jessen, C.A., Rundgren, M., Björk, S., Hammarlund D., 2005. Abrupt climatic changes and an
 730 unstable transition into a late Holocene Thermal Decline, a multiproxy lacustrine record from
 731 southern Sweden. J. Quat. Sci. 20, 349-362.

732

733 Jian, Z., Li, B., Plafman, U., Wang, P., 1996. Late Holocene cooling event in the western
 734 Pacific. Science in China Series D 39, 543-550.

735

736 Jiang, W.Y., Liu, T.S., 2007. Timing and spatial distribution of mid-Holocene drying over
 737 northern China, Response to a southeastward retreat of the East Asian Monsoon. J. Geophys.
 738 Res. 112, D24111, doi,10.1029/2007JD009050

739

740 Jin, G.Y., Liu, D.S., 2002. Mid-Holocene climate change in North China, and the effect on
 741 cultural development. Chin. Sci. Bull. 47, 408–413.

742

743 Lee, C-Y., Liew, P-M., Lee, T-Q.. 2010. Pollen records from southern Taiwan, implications for
 744 East Asian summer monsoon variation during the Holocene. Holocene 20, 81-89.

745

746 Li, C., Zheng, Y., Yu, S., Li, Y., Shen, H., 2012. Understanding the ecological background of
 747 rice agriculture on the Ningshao Plain during the Neolithic Age: pollen evidence from a buried
 748 paddy field at the Tianluoshan cultural site. *Quat. Sci. Rev.* 35, 131-138.

749

750 Li, H.M, Li, W.Y, Liu, J.L., 1995. Quaternary floras. In: Li, X., (Ed.), *Fossil Floras of China*
 751 *through the Geological Ages*. Guangdong Science and Technology Press, Guangzhou, pp 552–
 752 588.

753

754 Li, M., Ge, Q., Hao, Z.,Zheng, J., He, S., 2010a. Characteristics of cold-warm variation in the
 755 Hetao region and its surrounding areas in China during the past 5000 years. *Clim. Past* 6, 475-
 756 481.

757

758 Li, Y., Wu, J., Hou, S., Shi, C., Mo, D., Liu, B., Zhou, L., 2010b. Palaeoecological records of
 759 environmental change and cultural development from the Liangzhu and Qujialing archaeological
 760 sites in the middle and lower reaches of the Yangtze River. *Quat. Int.* 227, 29-37.

761

762 Li, Z., Sun, N., Dodson, J., Zhou, X., 2012. Human activity and its impact on the landscape at
 763 the Xishanping site in the western Loess Plateau during 4800-4300 cal yr BP based on the fossil
 764 charcoal record. *J. Archaeol. Sci.* 39, 3141-3147.

765

766 Liew, P.M., Lee, C.Y., Kuo, C.M., 2006. Holocene thermal optimal and climate variability of
 767 East Asian monsoon inferred from forest reconstruction of a subalpine pollen sequence, Taiwan.
 768 Earth Planet. Sci. Lett. 250, 596-605.
 769
 770 Lim, J., Fujiki, T., 2011. Vegetation and climate variability in East Asia driven by low-latitude
 771 oceanic forcing during the middle to late Holocene. Quat. Sci. Rev. 30, 2487-2497.
 772
 773 Liu, F., Feng, Z., 2012. A dramatic climatic transition at ~4000 cal. yr BP and its cultural
 774 responses in Chinese cultural domains. Holocene 22, 1181-1197.
 775
 776 Liu, F., Zhang, Y., Feng, Z., Hou, G., Zhou, Q., Zhang, H., 2010a. The impacts of climate
 777 change on the Neolithic cultures of Gansu-Qinghai region during the late Holocene
 778 Megathermal. J. Geogr. Sci. 20, 417-430.
 779
 780 Liu, H., Lin, Z., Qi, X., Zhang, M., Zhang, Z., Du, J., 2010b. Multiple analysis of variation of the
 781 East Asian Monsoon during the Holocene. Quat. Int. 213, 74-78.
 782
 783 Liu, H., Yin, Y., Zhu, J., Zhao, F., Wang, H., 2010c. How did the forest respond to Holocene
 784 climate drying at the forest-steppe ecotone in northern China? Quat. Int. 227, 46-52.
 785
 786 Liu, K-B., 1988. Quaternary history of the temperate forests of China. Quat. Sci. Rev. 7, 1-20.
 787

788 Liu, K-B., Qiu, H-L., 1994. Late-Holocene pollen records of vegetational changes in China,
789 climate or human disturbance. *Terrestr. Atmos. Ocean Sci.* 5, 393-410.
790

791 Liu, K-B., Sun, S., Jiang X.. 1992, Environmental change in the Yangtze River delta since
792 12,000 years B.P. *Quat. Res.* 38, 32–45.
793

794 Liu, T., Chen, Z., Sun, Q., et al. 2012. Migration of Neolithic settlements in the Dongting Lake
795 area of the middle Yangtze River basin, China, Lake-level and monsoon climate responses.
796 *Holocene* 22, 649-657.
797

798 Liu, Z., Harrison. S.P., Kutzbach J, Otto-Bliesner B. 2004. Global monsoons in the mid-
799 Holocene and oceanic feedback. *Clim. Dynam.* 22, 157–82.
800

801 Long, T., Qin, J., Atahan, P., Mooney, S., Taylor, D., 2014. Rising waters: new
802 geoarchaeological evidence of inundation and early agriculture from former settlement sites on
803 the southern Yangtze Delta, China. *Holocene* 24, 546-558.
804

805 Lu, TL-D., 2007. Mid-Holocene climate and cultural dynamics in eastern Central China. In:
806 Anderson DG, Maasch KA and Sandweiss DH (Eds.), *Climate Change and Cultural Dynamics,*
807 *A Global Perspective on Mid-Holocene Transition.* Academic Press, London, pp. 297-329.
808

809 Lutaenko, K.A., Zhushchikhovskaya, I.S., Mikishin, Y.A., Popov, A.N., 2007. In: Anderson,
810 D.G., Maasch, K.A., Sandweiss, D.H. (Eds.) *Climate Change and Cultural Dynamics: A Global*
811 *Perspective on Mid-Holocene Transitions*. Academic Press, London, pp. 331-406.

812

813 Ma, C., Zhu, C., Zheng, C., Wu, C., Guan, Y., Zhao, Z., Huang, L., Huang, R., 2008. High-
814 resolution geochemistry records of climate changes since late-glacial from Dajiuhu peat in
815 Shennongjia Mountains, central China. *Chin. Sci. Bull.* 53, 28-41.

816

817 Ma, C., Zhu, C., Zheng, C., Yin, Q., Zhao, Z., 2009. Climate changes in East China since the last
818 Late-glacial inferred from high-resolution mountain peat humification records. *Science in China*
819 *Series D, Earth Sciences* 52, 118-131.

820

821 Magny, M., Vannière, B., Zanchetta, G., Fouache, E., Touchais, G., Petrika, L., Coussot, C.,
822 Walter-Simonnet, A-V., Arnaud, F., 2009. Possible complexity of the climatic event around
823 4300-3800 cal. BP in the central and western Mediterranean. *Holocene* 19, 823-833.

824

825 Maher, B.A., 2008. Holocene variability of the East Asian Monsoon from Chinese cave records,
826 a re-assessment. *Holocene* 18, 861-866.

827

828 Maher, B.A., Thompson, R., 2012. Oxygen isotopes from Chinese caves, records not of monsoon
829 rainfall but of circulation regime. *J. Quat. Sci.* 27, 615-624.

830

831 Makohonienko, M., Kitagawa, H., Naruse, T., Nasu, H., Momohara, A., Okuno, M., Fujiki, T.,
832 Liu, X., Yasuda, Y., Yin, H., 2004. Late-Holocene natural and anthropogenic vegetation changes
833 in the Dongbei Pingyuan (Manchurian Plain), northeastern China. *Quat. Int.* 123-125, 71-88.
834

835 Marchant, R., Hooghiemstra, H., 2004. Rapid environmental change in African and South
836 American tropics around 4000 years before present, a review. *Earth Sci. Rev.* 66, 217–60.
837

838 Mischke, D., Zhang, C., 2010. Holocene cold events on the Tibetan plateau. *Glob. Planet.*
839 *Change* 72, 155-163.
840

841 Mo, D., Li, F., Li, S.C., 1996. A preliminary study on the palaeoenvironment of the middle
842 Holocene in the Hulu River area in Gansu Province and its effects on human activity. *Acta*
843 *Geogr. Sin.* 51, 59-69.
844

845 Mo, D., Zhao, Z., Xu, J., Li, M., 2011 Holocene environmental changes and the evolution of the
846 Neolithic cultures in China. In: Martini, I. P., & Chesworth, W., (Eds.), *Landscapes and Societies*
847 299-319. Springer, Netherlands.
848

849 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen Analysis*. Blackwell Scientific
850 Publications, Oxford, UK.
851

852 Morrill, C., Overpeck, J. T., Cole, J. E., 2003. A synthesis of abrupt changes in the Asian
853 summer monsoon since the last deglaciation. *Holocene* 13, 465-476.

854

855 Morrill, C., Overpeck, J.T., Cole, J.E., Liu, K-B., Shen, C., Tang, L., 2006 Holocene variations
856 in the Asian monsoon inferred from the geochemistry of lake sediments in central Tibet. *Quat.*
857 *Res.* 65, 232-243.

858

859 Okuda, M., Sato, Y., Tang, L.H., Takahashi, M., Toyama, S., Yano, A., Kitagawa, H., Yasuda,
860 Y., 2003 Late Holocene vegetation and environment at Cauduntou, west of Yangtze Delta, SW
861 Jiangsu Province, East China. *Quat. Int.* 105, 39-47.

862

863 O’Sullivan, P., 2008. The ‘collapse’ of civilizations, what palaeoenvironmental reconstruction
864 cannot tell us, but anthropology can. *Holocene* 18, 45-55.

865

866 Olsen, J., Anderson, N.J., Knudsen, M.F., 2012. Variability of the North Atlantic Oscillation
867 over the past 5200 years. *Nature Geoscience* 5, 808–812.

868

869 Pals, J.P., van Geel, B., Delfos, A., 1980. Palaeoecological studies in the Klokkeweel bog near
870 Hoogkarspel (Noord Holland). *Rev. Palaeobot. Palynol.* 30, 371-418.

871

872 Peiser, B.J., 1998. Comparative analysis of late Holocene environmental and social upheaval,
873 evidence for a disaster around 4000 BP. In: *Natural Catastrophes during Bronze Age*
874 *Civilisations*. Peiser, B.J., Palmer, T., Bailey, M., (Eds.), British Archaeological Reports,
875 International Series 728, 117–139.

876

877 Perry, C.A., Hsu, K.J., 2000. Geophysical, archaeological, and historical evidence support a
 878 solar-output model for climate change. *Proceedings of the National Academy of Sciences* 97,
 879 12,433–12,438.
 880
 881 Phadtare, N.R., 2000. Sharp decrease in Summer Monsoon strength 4000-3500 cal yr B.P. in the
 882 central higher Himalaya of India based on pollen evidence from alpine peat. *Quat. Res.* 53, 122-
 883 129.
 884
 885 Qin, J., Taylor, D., Atahan, P., Zhang, X., Wu, G., Dodson, J., Zheng, H., Itzstein-Davey, F.,
 886 2011. Neolithic agriculture, freshwater resources and rapid environmental changes on the lower
 887 Yangtze, China. *Quat. Res.* 75, 55-65.
 888
 889 Qu, W., Xue, B., Dickman, M.D., Wang, S., Fan, C., Wu, R., Zhang, P., Chen, J., Wu, Y., 2000.
 890 A 14 000 year record of palaeoenvironmental change in the western basin of China's third
 891 largest lake, Lake Taihu. *Hydrobiol.* 432, 113-120.
 892
 893 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey,
 894 C., Buck, C.E., Burr, G.S., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P.,
 895 Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac, F.G.,
 896 Manning, S.W., Reimer, R.W., Richards, D.A., Southon, J.R., Talamo, S., Turney, C.S.M., van
 897 der Plicht, J., Weyhenmeyer, C.E., 2009. IntCal09 and marine09 radiocarbon age calibration
 898 curves, 0-50,000 years cal BP. *Radiocarbon* 51, 1111-1150.
 899

900 Ren, G., 2000. Decline of the mid- to late Holocene forests in China, climatic change or human
 901 impact? *J. Quat. Sci.* 15, 273–281.
 902
 903 Ren, G., 2007. Changes in forest cover in China during the Holocene. *Veget. Hist. Archaeobot.*
 904 16, 119-126.
 905
 906 Ren, G., Beug, H-J., 2002. Mapping Holocene pollen data and vegetation of China. *Quat. Sci.*
 907 *Rev.* 21, 1395-1422.
 908
 909 Ren, G., Zhang, L., 1998. A preliminary mapped summary of Holocene pollen data for Northeast
 910 China. *Quat. Sci. Rev.* 17, 669–688.
 911
 912 Renssen, H., Seppä, H., Crosta, X., Goosse, H., Roche, D.M., 2012. Global characterization of
 913 the Holocene thermal maximum. *Quat. Sci. Rev.* 48, 7-19.
 914
 915 Riehl, S., 2012. Variability in ancient Near Eastern environmental and agricultural development
 916 *J. Arid Env.* 86, 113-121.
 917
 918 Sandweiss D.H., Maasch, K.A., Anderson D.G., 1999. Transitions in the mid-Holocene. *Science*
 919 283, 499-500.
 920

921 Schettler, G., Liu, Q., Mingram, J., Stebich, M., Dulski, P., 2006. East-Asian Monsoon
 922 variability between 15 000 and 2000 cal. yr BP recorded in varved sediments of Lake
 923 Sihailongwan (northeastern China, Long Gang volcanic field). *Holocene* 16, 1043-1057.
 924
 925 Schofield, J.E., Waller, M.P., 2005. A pollen analytical record for hemp retting from Dungeness
 926 Foreland, UK. *J. Archaeol. Sci.* 32, 715-726.
 927
 928 Schulmeister, J., Lees, B.G., 1995. Pollen evidence from tropical Australia for the onset of an
 929 ENSO-dominated climate at c. 4000 BP. *Holocene* 5, 10-18.
 930
 931 Selvaraj, K., Chen, C.T.A., Lou, J-H., 2007. Holocene East Asian monsoon variability, links to
 932 solar and tropical Pacific forcing. *Geophys. Res. Lett.* 34, L01703.
 933
 934 Shao, X., Wang, Y., Cheng, H., Kong, X., Wu, J., Edwards, R.L., 2006. Long-term trend and
 935 abrupt events of the Holocene Asian monsoon inferred from a stalagmite $d^{18}O$ record from
 936 Shennongjia in central China. *Chin. Sci. Bull.* 51, 1–8.
 937
 938 Shi, W., Ma, C.M., Zhu, C., Wang, F.B., Li, S.J., 2008. Analysis of stratigraphy on multi-profiles
 939 in the Taihu Lake region and palaeoenvironmental events in the Liangzhu culture epoch. *Geogr.*
 940 *Res.* 27, 1129-1138 (in Chinese).
 941
 942 Shi, Y., Kong, Z., Wang, S., Tang, L., Wang, F., Yao, T., Zhao, X., Zhang, P., Shi, S., 1993.
 943 Mid-Holocene climates and environments in China. *Glob. Planet. Change* 7, 219–233.

944

945 Shi, Y., Kong, Z., Wang, S., Tang, L., Wang, F., Yao, T., Zhao, X., Zhang, P., Shi, S., 1994.

946 Climates and environments of the Holocene Megathermal Maximum in China. *Science in China*

947 (Series B), 37, 481-493.

948

949 Shu, J-W., Wang, W-M., Chen, W., 2007. Holocene vegetation and environment changes in the

950 NW Taihu plain, Jiangsu Province, East China. *Acta Micropalaeontol. Sin.* 24, 210-221. (in

951 Chinese).

952 Stanley, D.J., Chen, Z., Song, J., 1999. Inundation, sea-level rise and transition from Neolithic to

953 Bronze Age cultures, Yangtze Delta, China. *Geoarchaeology* 14, 15-26.

954 Staubwasser, M., Weiss, H., 2006. Holocene climate and cultural evolution in late prehistoric-

955 early historic West Asia. *Quat. Res.* 66, 372-87.

956 Staubwasser, M., Sirocko, F., Grootes, P.M., Segl, M., 2003. Climate change at 4.2 ka BP

957 termination of the Indus Valley civilization and Holocene South Asian monsoon variability.

958 *Geophys. Res. Lett.* 30, 1425-1428.

959

960 Stott, L., Cannariato, K., Thunell, R., Haug, G.H., Koutavas, A., Lund, S., 2004. Decline of

961 surface temperature and salinity in the western tropical Pacific Ocean in the Holocene epoch.

962 *Nature* 431, 56-59.

963

964 Sun, D., Gagan, M.K., Cheng, H., Scott-Gagan, H., Dykoski, C., Edwards, R.L., Su, R., 2005.

965 Seasonal and interannual variability of the mid-Holocene East Asian monsoon in coral $\delta^{18}\text{C}$

966 records from the South China Sea. *Earth Planet. Sci. Lett.* 237, 69-84.

967

968 Sun, X., Chen, Y., 1991. Palynological records of the last 11,000 years in China. *Quat. Sci. Rev.*

969 10, 537–544.

970

971 Sun, X., Weng, C., 1992. Pollen records on the history of mixed conifer and hardwood forest in

972 Northeast China. *Acta Bot. Sin.* 34, 394–401 (in Chinese).

973

974 Sun, X., Luo, Y., Huang, F., Tian, J., Wang, P., 2003. Deep-sea pollen from the South China

975 Sea: Pleistocene indicators of East Asian monsoon. *Mar. Geol.* 201, 97-118.

976

977 Tan, M., Hou, J., Liu, T., 2004. Sun-coupled climate connections between Eastern Asia and

978 Northern Atlantic. *Geophys. Res. Lett.* 31, 7207.

979

980 Tan, L.C., An, Z.S., Cai, Y.J., Long, H., 2008. Hydrological representation of the 4.2 ka BP

981 event in China and its global linkages. *Geol. Rev.* 54, 94-104 (in Chinese).

982

983 Tao, J., Chen, M-T., Xu, S., 2006. A Holocene environmental record from the southern Yangtze

984 River delta, eastern China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 230, 204-229.

985

986 van Geel, B., 1986. Application of fungal and algal remains and other microfossils in

987 palynological analyses. In: Berglund, B.E., (Ed.), *Handbook of Palaeoecology and Palynology.*

988 John Wiley, New York, pp. 497-505.

989

990 van Geel, B., 2001. Non-pollen palynomorphs. In: Smol, J.P., Birks H.J.B., Last, W.M., (Eds.),
 991 Tracking Environmental Change using Lake Sediments. Volume 3, Terrestrial, Algal and
 992 Siliceous Indicators. Kluwer, Dordrecht, pp. 99-119.
 993
 994 van Geel, B., Aptroot, A., 2006. Fossil ascomycetes in Quaternary deposits. Nova Hedwigia 82,
 995 313-329.
 996
 997 van Geel, B., Mur, L.R., Ralska- Jasiewiczowa, M., Goslar, T., 1994. Fossil akinetes of
 998 *Aphanizomenon* and *Anabaena* as indicators of medieval phosphate-eutrophication of Lake
 999 Gosciarz (Central Poland). Rev. Palaeobot. Palynol. 83, 97-105.
 1000
 1001 van Geel, B., Odgaard, B.V., Ralska-Jasiewiczowa, M., 1996. Cyanobacteria as indicators of
 1002 phosphate-eutrophication of lakes and pools in the past. PACT 50, 399-415.
 1003
 1004 Walker, M.J.C., Berkelhammer, M., Björk, S., Cwynar, L.C., Fisher, D.A., Long, A.J., Lowe,
 1005 J.J., Newnham, R.M., Rasmussen, S.O., Weiss, H., 2012. Formal subdivision of the Holocene
 1006 Series/Epoch, a discussion paper by a working group of INTIMATE (Integration of ice-
 1007 core,marine and terrestrial records) and the subcommission on Quaternary stratigraphy
 1008 (International commission on Stratigraphy). J. Quat. Sci. 27, 649-659.
 1009
 1010 Wang, B., Wu, R., Fu, X., 2000. Pacific-east Asian teleconnection, how does ENSO affect east
 1011 Asian climate? J. Clim. 13, 1517-1536.
 1012

1013 Wang, F., Chien, N., Zhang, Y., Yang, H., 1995. Pollen Flora of China (second edition). Science
1014 Press, Beijing (in Chinese).

1015

1016 Wang, J., Chen, X., Zhu, X.H., Liu, J.L., Chang, W.Y.B., 2001. Taihu Lake, lower Yangtze
1017 drainage basin, evolution, sedimentation rate and the sea level. *Geomorphol.* 41, 183–193.

1018

1019 Wang, L., Sarnthein, M., Erlenkeuser, H., Grootes, P.M., Grimalt, J.O., Pelejero, C., Linck, G.,
1020 1999a. Holocene variations in Asian Monsoon moisture: a bidecadal sediment record from the
1021 South China Sea. *Geophys. Res. Lett.* 26, 2889-2892.

1022

1023 Wang, L., Sarnthein, M., Erlenkeuser, H., Grimalt, J.O., Grootes, P.M., Heilig, S., Ivanova, E.,
1024 Kienast, M., Pelejero, C., Pflaumann, U., 1999b. *Mar. Geol.* 156, 245-284.

1025

1026 Wang, S., Lü, H., Liu, J., Negendank, J.F.W., 2007. The early Holocene optimum inferred from
1027 a high-resolution pollen record of Huguangyan Maar Lake in southern China. *Chin. Sci. Bull.* 52,
1028 2829-2836.

1029

1030 Wang, Y.L., Cheng, H., Edwards, R.L., An, Z.S., Wu, J.Y., Shen, C.C., Dorale, J.A., 2001. A
1031 high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China. *Science*
1032 294, 2345–2348.

1033 Wang, Y., Cheng, H., Edwards, R.L., He, Y., Kong, X., An, Z., Wu, J., Kelly, M.J., Dykoski,
1034 C.A., Li, X., 2005. The Holocene Asian monsoon, links to solar changes and North Atlantic
1035 climate. *Science* 308, 854-857.

1036

1037 Wang, Y., Liu, X., Herzschuh, U., 2010. Asynchronous evolution of the Indian and East Asian
 1038 summer monsoon indicated by Holocene moisture patterns in monsoonal central Asia. *Earth-Sci.*
 1039 *Rev.* 103, 135-153.

1040

1041 Wang, Z., Li, M., Zhang, R., Zhuang, C., Liu, Y., Saito, Y., Xie, J., Zhao, B., 2011. Impacts of
 1042 human activity on the late-Holocene development of the subaqueous Yangtze Delta, China, as
 1043 shown by magnetic properties and sediment accumulation rates. *Holocene* 21, 393-407.

1044

1045 Wang, Z., Zhuang, C., Saito, Y., Chen, J., Zhan, Q., Wang, X., 2012. Early mid-Holocene sea-
 1046 level change and coastal environmental response on the southern Yangtze delta plain, China,
 1047 implications for the rise of Neolithic culture. *Quat. Sci. Rev.* 35, 51-62.

1048

1049 Wanner H, Solomina O, Grosjean M, Ritz, S.P., Jetel, M., 2011. Structure and origin of
 1050 Holocene cold events. *Quat. Sci. Rev.* 30, 3109-3123.

1051

1052 Wei, K., Lee, M., Duan, W., Chen, C., Wang, C., 1998. Palaeoceanographic change in the
 1053 northeastern South China Sea during the last 15000 years. *J. Quat. Sci.* 13, 55-64.

1054

1055 Wei, K., Gasse, F., 1999. Oxygen isotopes in lacustrine carbonates of West China revisited,
 1056 implications for post glacial changes in summer monsoon circulation. *Quat. Sci. Rev.* 18, 1315-
 1057 1334.

1058

1059 Weiss, H., Courty, M.A., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R., Curnow, A.,
 1060 1993. The genesis and collapse of Third Millennium north Mesopotamian civilization. *Science*
 1061 261, 995–1004.
 1062
 1063 Wen, R., Xiao, J., Chang, Z., Zhai, D., Xu, Q., Li, Y., Itoh, S., Lomtatidze, Z., 2010. Holocene
 1064 climate changes in the mid-high-latitude-monsoon margin reflected by the pollen record from
 1065 Hulun Lake, northeastern Inner Mongolia. *Quat. Res.* 73, 293-303.
 1066
 1067 Wen, R., Xiao, J., Ma, Y., Feng, Z., Li, Y., Xu, Q., 2013. Pollen-climate transfer functions
 1068 intended for temperate eastern Asia. *Quat. Int.* 311, 3-11.
 1069
 1070 Wen, R., Xiao, J., Chang, Z., Zhai, D., Xu, Q., Li, Y., Itoh, S., 2010. Holocene precipitation and
 1071 temperature variations in the East Asian monsoonal margin from pollen data from Hulun Lake in
 1072 northeastern Inner Mongolia, China. *Boreas* 39, 262-272.
 1073
 1074 Wu, L., Wang, X.Y., Zhou, K.S., Mo, D., Zhu, C., Gao, C., Zhang, G., Li, L., Han, W., 2010.
 1075 Transmutations of ancient settlements and environmental changes between 6000 and 2000 aBP
 1076 in the Chaohu Lake Basin, East China. *J. Geogr. Sci.* 20, 687-700.
 1077
 1078 Wu, L., Li, F., Zhu, C., Li, B., 2012. Holocene environmental change and archaeology, Yangtze
 1079 River Valley, China, review and prospects. *Geosci. Frontiers* 3, 875-892.
 1080

1081 Wu, L., Zhu, C., Zheng, C., Li, F., Wang, X., Li, L., Sun, W., 2014. Holocene environmental
 1082 change and its impacts on human settlement in the Shanghai area, east China. *Catena* 114, 78-89.
 1083
 1084 Wu, W., Liu, T., 2004. Possible role of the “Holocene Event 3” on the collapse of Neolithic
 1085 Cultures around the Central Plain of China. *Quat. Int.* 117, 153-166.
 1086
 1087 Wu X, Zhang Z, Xu X, Shen J., 2012. Asian summer monsoonal variations during the Holocene
 1088 revealed by Huguangyan maar lake sediment record. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*
 1089 323–325, 13–21.
 1090
 1091 Xia, Z.K., Yang, X., Ye, M.L., 2003. Prehistoric disasters at Lajia site, Qinghai, China. *Chin.*
 1092 *Sci. Bull.* 48, 1877- 1881.
 1093
 1094 Xiao, J., Xu, Q.H., Nakamura, T, Yang, X., Liang, W., Inouchi, Y., 2004. Holocene vegetation
 1095 variation in the Daihai Lake region of north-central China, a direct indication of the Asian
 1096 monsoon climatic history. *Quat. Sci. Rev.* 23, 1669-1679.
 1097
 1098 Xu, D., Lu, H., Wu, N., Liu, Z., 2010. 30 000-year vegetation and climate change around the
 1099 East China Sea shelf inferred from a high resolution pollen record. *Quat. Int.* 227, 53-60.
 1100 Xu, X., Chang, W.Y.B., Liu, J., 1996. Changes in vegetation and climate in the Taihu Lake
 1101 during the last 11,000 years. *Acta Palaeontol. Sin.* 35, 175-186. (in Chinese).
 1102

1103 Xu. Q.H., Xiao, J.L., et al. 2010. Pollen-based quantitative reconstruction of Holocene climate
 1104 changes in the Daihai Lake Area, Inner Mongolia, China. *J. Clim.* 23, 2856–2868.
 1105
 1106 Yan, Q., Xu, S., Shao, X., 1989. Holocene cheniers in the Yangtze delta, China. *Mar. Geol.* 90,
 1107 337–343.
 1108
 1109 Yancheva, G., Nowaczyk, N.R., Mingram, J., Dulski, P., Schettler, G., Negendank, J.F.W., Liu,
 1110 J., Sigman, D.M., Peterson, L.C., Haug, G.H., 2007. Influence of the intertropical convergence
 1111 zone on the East Asian monsoon. *Nature* 445: 74-77.
 1112
 1113 Yang, X., Scuderi, L.A., 2010. Hydrological and climatic changes in deserts of China since the
 1114 late Pleistocene. *Quat. Res.* 73, 1-9.
 1115
 1116 Yao, P., Huang, C.C., Pang, J.L., Zha, X.C., Li, X.G., 2008. Palaeoflood hydrological studies in
 1117 the middle reaches of the Beiluohe River. *Acta Geogr. Sin.* 63, 1198-1206 (in Chinese).
 1118
 1119 Yasuda, Y., Fujiki, T., Nasu, H., Kato, M., Morita, Y., Mori, Y., Kanehara, M., Toyama, S.,
 1120 Yano, A., Okuno, M., Jiejun, H., Ishihara, S., Kitagawa, H., Fukusawa, H., Naruse, T., 2004.
 1121 Environmental archaeology at the Chengtoushan site, Hunan Province, China, and implications
 1122 for environmental change and the rise and fall of the Yangtze river civilization. *Quat. Int.* 123-
 1123 125, 149-158.
 1124

1125 Yi, S., Saito, Y., Zhao, Q., Wang, P., 2003. Vegetation and climate changes in the Changjiang
 1126 (Yangtze River) Delta, China, during the past 13,000 years inferred from pollen records. *Quat.*
 1127 *Sci. Rev.* 22, 1501-1519.
 1128
 1129 Yi, S., Saito, Y., Yang, D-Y., 2006. Palynological evidence for Holocene environmental change
 1130 in the Changjiang (Yangtze River) delta, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 241,
 1131 103–117.
 1132
 1133 Yu, S., Zhu, C., Song, H., Qu, W., 2000. Role of climate in the rise and fall of Neolithic cultures
 1134 on the Yangtze Delta. *Boreas* 29, 157-165.
 1135
 1136 Yu, S., Zhu, C., Wang, F., 2003. Radiocarbon constraints on the Holocene flood deposits of the
 1137 Ning-Zhen mountains, lower Yangtze River area of China. *J. Quat. Sci.* 18, 521-525.
 1138
 1139 Yu, X., Zhou, W., Franzen, L.G., Xian, F., Cheng, P., Jull, A.J.T., 2006. High-resolution peat
 1140 records for Holocene monsoon history in the eastern Tibetan plateau. *Science in China Series D,*
 1141 *Earth Sciences* 49, 615-621.
 1142
 1143 Yuan, D., Cheng, H., Edwards, R.L., Dykoski, C.A., Kelly, M.J., Zhang, M., Qing, J., Lin, Y.,
 1144 Wang, Y., Wu, J., Dorale, J.A., An, Z., Cai, Y., 2004. Timing, duration and transitions of the last
 1145 interglacial Asian monsoon. *Science* 304, 575–578.
 1146

1147 Zhai, D., Xiao, J., Zhou L., Wen, R., Chang, Z., Wang, X., Jin, X., 2011. Holocene East Asian
 1148 Monsoon variation inferred from species assemblage and shell chemistry of the ostracodes from
 1149 Hulun Lake, Inner Mongolia. *Quat. Res.* 75, 512-522.
 1150
 1151 Zhang, D.D., Lee, H.F., Wang, C., Li, B., Zhang, J., Pei, Q., Chen, J., 2011. Climate change and
 1152 large-scale human population collapses in the pre-industrial era. *Glob. Ecol. Biogeogr.* 20, 520-
 1153 531.
 1154
 1155 Zhang, G., Zhu, C., Wang, J., Zhu, G., Ma, C., Zheng, C., Zhao, L., Li, Z., Li, L., Jin, A., 2010.
 1156 Environmental archaeology on Longshan culture (4500-4000 a BP) at Yuhuicun site in Bengbu,
 1157 Anhui Province. *J. Geogr. Sci.* 20 (3), 455-468.
 1158
 1159 Zhang, J., Chen, F., Holmes, J.A., Li, H., Guo, X., Wang, J., Li, S., Lü Y., Zhao, Y., Qiang, M.,
 1160 2011. Holocene monsoon climate documented by oxygen and carbon isotopes from lake
 1161 sediments and peat bogs in China, a review and synthesis. *Quat. Sci. Rev.* 30, 1973-1987.
 1162
 1163 Zhang, Q., Jiang, T., Shi ,Y., King, L., Liu, C., Metzler, M., 2004a. Palaeo-environmental
 1164 changes in the Yangtze Delta during past 8000 years. *J. Geogr. Sci.* 14, 105-112.
 1165
 1166 Zhang, Q., Liu, C., Zhu, C., Jiang, T., 2004b. Environmental change and its impacts on human
 1167 settlement in the Changjiang river delta in Neolithic age. *Chin. Geogr. Sci.* 14, 239-244.
 1168

1169 Zhang, Q., Zhu, C., Liu, C.L., Jiang, T., 2005. Environmental change and its impacts on human
 1170 settlement in the Yangtze Delta, P.R. China. *Catena* 60, 267-277.
 1171
 1172 Zhang, Y., Xi, Y., Zhang, J., Gao, G., Du, N., Sun, X., Kong, Z., 1990. Spore Morphology of
 1173 Chinese Pteridophytes. Science Press, Beijing.
 1174
 1175 Zhang, Y., 2007. Causes of Liangzhu culture's disappearance from micropalaeontology. *Sci.*
 1176 *Geogr. Sin.* 27, 376-379.
 1177
 1178 Zhao, Y., Hölzer, A., Yu, Z., 2007. Late Holocene natural and human-induced environmental
 1179 change reconstructed from peat records in eastern central China. *Radiocarbon* 49, 789-798.
 1180
 1181 Zhao, Y., Yu, Z., Chen, F., Zhang, J., Yang, B., 2009. Vegetation response to Holocene climate
 1182 change in monsoon-influenced region of China. *Earth-Sci. Rev.* 97, 242-256.
 1183
 1184 Zheng, Y.Q., Yu, G., Wan, S.M., Xue, B., Zhuo, D.Q., Zeng, X.M., Liu, H.Q., 2004. Simulation
 1185 of paleoclimate over East Asia at 6 ka BP and 21 ka BP by a regional climate model. *Clim.*
 1186 *Dynam.* 23, 513-529.
 1187
 1188 Zheng, Y., Sun, G., Qin, L., Li, C., Wu, X., Chen, X. 2009. Rice fields and modes of rice
 1189 cultivation between 5000 and 2500 BC in East China. *J. Archaeol. Sci.* 36, 2609-2616.
 1190

1191 Zheng, Z., Li, Q., 2000. Vegetation, climate and sea level in the past 55,000 years, Hanjiang
1192 delta, southeastern China. *Quat. Res.* 53, 330-340.
1193

1194 Zhong, W., Xue, J., Zheng, Y., Ouyang, J., Ma, Q., Cai, Y., Tang, X. 2010. Climatic changes
1195 since the last deglaciation inferred from a lacustrine sedimentary sequence in the eastern Nanling
1196 Mountains, south China. *J. Quat. Sci.* 25, 975-984.
1197

1198 Zhou, X., Li, X., 2012. Variations in spruce (*Picea* sp.) distribution in the Chinese Loess Plateau
1199 and surrounding areas during the Holocene. *Holocene* 22, 687-696.
1200

1201 Zhu, C., Zhang, Q., 2006. Climatic evolution in the Yangtze Delta region in the late Holocene
1202 epoch. *J. Geogr. Sci.* 16, 423-429.
1203

1204 Zhu, C., Song, J., You, K., Han, H., 1996. Research on the reasons of culture interruption in
1205 Maqiao Site, *Chin. Sci. Bull.* 41, 148-152. (in Chinese).
1206

1207 Zhu, C., Ma, C., Yu, S-Y., Tang, L., Zhang, W., Lu, X. 2010. A detailed pollen record of
1208 vegetation and climate changes in Central China during the past 16 000 years. *Boreas* 39, 69-76.
1209

1210 Zhuang, Y., Ding, P., French, C., 2014. Water management and agricultural intensification of
1211 rice farming at the late-Neolithic site of Maoshan, Lower Yangtze River, China. *Holocene* 24,
1212 531-545.
1213

1214 Zong, Y., Chen, Z., Innes, I.B., Chen, C., Wang, Z., Wang, H. 2007. Fire and flood management
 1215 of coastal swamp enabled first rice paddy cultivation in east China. *Nature* 449, 459-462.
 1216

1217 Zong, Y., Innes, J.B., Wang, Z., Chen, Z., 2011. Mid-Holocene coastal hydrology and salinity
 1218 changes in the east Taihu area of the lower Yangtze wetlands, China. *Quat. Res.* 76, 69-82.
 1219

1220 Zong, Y., Wang, Z., Innes, J.B., Chen, Z., 2012a. Holocene environmental change and Neolithic
 1221 rice agriculture in the lower Yangtze region of China: a review. *Holocene* 22, 623-635.
 1222

1223 Zong, Y., Innes, J.B., Wang, Z., Chen, Z., 2012b. Environmental change and Neolithic
 1224 settlement movement in the lower Yangtze wetlands of China. *Holocene* 22, 659-673.
 1225

1226 Zou, S., Cheng, G., Xiao, H., Xu, B., Feng, Z., 2009. Holocene natural rhythms of vegetation and
 1227 present potential ecology in the Western Chinese Loess Plateau. *Quat. Int.* 194, 55-67.
 1228
 1229
 1230
 1231
 1232
 1233
 1234
 1235
 1236

1237 Table 1. Lithostratigraphy at Pingwang (30° 57' 30"N; 120° 38' 25"E; altitude 1.6 m YSD)

Depth (m)	Descriptions
0.00 – 0.45	Paddy field soils
0.45 – 1.10	Brown to yellowish grey, hard to firm, clay
1.10 – 1.25	Blackish grey, soft, clay
1.25 – 1.80	Greenish grey, soft, clay
1.80 – 2.10	Brownish grey, soft, organic rich clay
2.10 – 3.10	Dark grey, soft, organic rich clay with small shells found in upper part
3.10 – 3.70	Greenish grey, soft, silt and clay
3.70 – 3.80	Blackish brown peat
3.80 – 4.00	Sticky hard clay (pre-Holocene)

1238
1239 Table 2. AMS Radiocarbon dates at Pingwang. Calibrated results are shown, using the Calib6.1
1240 and IntCal09 programmes (Reimer et al., 2009). There is an inversion in the bottom two dates,
1241 but based on regional pollen and radiocarbon data Beta-253340 is more likely to be correct.

1242

Depth (m)	Dated material	¹⁴ C date (a BP)	Calibrated age range (cal BP) (2σ)	Mid-point age (cal BP)	Laboratory code
1.59-1.61	pollen residue	2700±40	2750-2869	2810±60	Beta-255432
1.85-1.87	pollen residue	4430±40	4872-5280	5076±204	Beta-266433
2.25-2.27	plant macrofossil	4720±40	5324-5584	5454±130	Beta-243208
3.20-3.22	pollen residue	6800±50	7573-7724	7649.5±75.5	Beta-253340
3.75-3.77	peat	6290±50	7026-7322	7174±148	Beta-228442

Captions to figures


Fig. 1. Location of the study area, around Taihu lake in the Yangtze coastal plain, showing the topography and the position of other, smaller lakes, the barrier ridges to the east that protect the plain from the sea, and the location of the coring site at Pingwang. Figures 1a and 1b show the distribution of the main Late Neolithic archaeological sites of the Liangzhu and Maqiao cultures.

Fig. 2. Percentage tree and shrub pollen diagram from Pingwang. Frequencies are calculated as percentages of the total tree and shrub sum, with major stages in vegetation history and summary curves for ecological groupings also shown. Calibrated radiocarbon age ranges before present (cal BP) are shown on the left of the diagram, which is zoned at major changes in the tree and shrub pollen curves using the CONISS program. The timings of the following major environmental events are shown on the diagram. 1: Early in the mid-Holocene Megathermal 2: The end of major saline estuarine influence at the site (after Zong et al., 2011) 3: The maximum of the Megathermal optimum 4: The start of the climatic decline towards the Late Holocene 5: Abrupt climate deterioration ca 4200 cal BP. Depths are in centimetres below present ground surface.


Fig. 3. Percentage herb pollen and pteridophyte spore diagram from Pingwang, frequencies calculated as percentages of the total tree and shrub pollen sum. Diagram zonation, calibrated radiocarbon dates and environmental events follow Fig. 2. Depths are in centimetres below present ground surface.

1268 Fig. 4 Percentage non-pollen palynomorph (NPP) diagram from Pingwang. NPPs and
1269 microscopic charcoal fragment (microcharcoal) frequencies are shown as percentages of the tree
1270 and shrub pollen sum. X axis NPP type numbers refer to the catalogue of the Hugo de Vries
1271 laboratory, Amsterdam, and where taxon names are unknown they are assigned the prefix HdV.
1272 Diagram zonation, radiocarbon dates and environmental events follow Fig. 2. Depths are in
1273 centimetres below present ground surface.
1274

*Figure



*Figure



*Figure

